

UHPLC Analysis of Vitamin D2 & D3 Metabolites and Epimers on Supel™ Carbon LC Column

Introduction

Analysis of vitamin D metabolites has continued to be a topic of interest in recent publications, primarily as biomarkers for possible disease states and vitamin deficiency. While vitamin D is present in two forms, vitamin D3 and vitamin D2, current ELISA methods demonstrate different cross-reactivities and cannot distinguish between D2 and D3 forms of the vitamin metabolites resulting in erroneous reporting of total

25-hydroxyvitamin D concentrations (**Figure 1**). Further, there is interest in an analytical means to differentiate the D2 and D3 forms from the D2 and D3 epimers because of their different degrees of bioactivity. This application demonstrates the use of the Supel $^{\text{TM}}$ Carbon LC UHPLC column, with its ability to resolve structural isomers, to baseline separate all four analytes.

25-hydroxyvitamin D3

. .

3-epi-25-hydroxyvitamin D3

3-epi-25-hydroxyvitamin D2

Figure 1: Structures of Vitamin D2 and D3 metabolites.

25-hydroxyvitamin D2



Results/Conclusion

Figure 2 shows the results of the separation of the Vitamin D2 and D3 metabolites. Note the sharp peak shape and baseline resolution between all four analytes.

Vitamin D2 & D3 Separation on Supel™ Carbon LC column




Figure 2: Chromatographic separation of Vitamin D2 and D3 metabolites on Supel™ Carbon LC. Conditions: Column: Supel™ Carbon LC, 10 cm x 2.1 mm I.D., 2.7 μm; Mobile Phase: [A] 2-Propanol; [B] Tetrahydrofuran; Gradient: 0% B to 70% B in 15 min; hold at 70% B for 5 min; Flow Rate: 0.3 mL/min; Column Temp.: 25 °C; Detector: UV, 275 nm; Injection: 2.0 μL; Sample: Vitamin D2 and D3 metabolites mix, varied concentration, ethanol

This application demonstrated the use of the Supel™ Carbon LC column to resolve vitamins D2 and D3 metabolites and their epimers. Baseline separation of all four analytes was achieved with excellent peak shape and sensitivity.

Product list	PN
Supel™ Carbon LC Column, 10 cm x 2.1mm I.D., 2.7 μm	59986-U
2-Propanol for HPLC, 99.5%	439207
Tetrahydrofuran for HPLC, > 99.9%, Inhibitor-free	439215
25-hydroxyvitamin D3 solution, 100 μg/mL in ethanol	H-083
25-hydroxyvitamin D2 solution, 50 μg/mL in ethanol	H-073
3-epi-25-hydroxyvitamin D3, 50 μg/mL in ethanol	E-086
3-epi-25-hydroxyvitamin D2, 98%	753149

MilliporeSigma 400 Summit Drive Burlington, MA 01803

To place an order or receive technical assistance

Order/Customer Service: SigmaAldrich.com/order Technical Service: SigmaAldrich.com/techservice Safety-related Information: SigmaAldrich.com/safetycenter

SigmaAldrich.com

